

Occupation number and fluctuations in the finite-temperature Bose-Hubbard model

L. I. Plimak and M. Fleischhauer

Fachbereich Physik, Technische Universität Kaiserslautern, D-67663 Kaiserslautern, Germany.

M. K. Olsen

*Instituto de Física da Universidade Federal Fluminense,
Boa Viagem 24210-340, Niterói, Rio de Janeiro, Brazil.*

(Dated: February 2, 2008)

We study the occupation numbers and number fluctuations of ultra-cold atoms in deep optical lattices for finite temperatures within the Bose-Hubbard model. Simple analytical expressions for the mean occupation number and number fluctuations are obtained in the weak-hopping regime using an interpolation between results from different perturbation approaches in the Mott-insulator and superfluid phases. These analytical results are compared to exact one dimensional numerical calculations using a finite temperature variant of the Density-Matrix Renormalisation Group (DMRG) method and found to have a high degree of accuracy. We also find very good agreement in the crossover “thermal” region. With the present approach the magnitude of number fluctuations under realistic experimental conditions can be estimated and the properties of the finite temperature phase diagram can be studied.

PACS numbers: 39.25.+k, 03.75.Mn, 05.30.JP

The Bose-Hubbard (BH) model, first studied in the late 1980’s [1], has long been considered a rather academic testing ground for analytical as well as numerical approaches in quantum statistical mechanics. Its most interesting feature is the existence of a quantum phase transition [2] at zero temperature between a superfluid and a Mott-insulating phase. With recent advances in the experimental techniques of atom trapping and optical lattices, the BH model has regained substantial practical importance. As first noticed in [3], the BH Hamiltonian applies to bosonic atoms trapped in a deep lattice potential. Recent experiments with optical lattices [4] have spectacularly confirmed the existence of the superfluid-insulator transition. Since the Mott-phase is characterized by a well defined occupation number of the potential wells, the phase transition has important potential applications in quantum information processing [5, 6], or Heisenberg-limited matter-wave interferometry [7], both of which require optical lattices with regular filling. Although techniques to eliminate lattice defects have been designed [8] an important issue for the implementation of these applications under experimental conditions are fluctuations in the particle number per site at finite T . In this work we analyze these fluctuations by studying the finite temperature Bose-Hubbard model.

The zero temperature BH model has been extensively investigated in the past by methods such as the Gutzwiller projection ansatz [9], the strong-coupling expansion [10, 11, 12, 13] and Quantum Monte-Carlo [14, 15, 16, 17, 18], as well as with various mean-field approaches (see, e.g., [1, 19, 20]). A very powerful numerical technique in the case of one spatial dimension is the density matrix renormalization group (DMRG) [21], used to calculate first and second order (i.e., amplitude and number-number) correlations [22]. Less attention has been paid to the nonzero temperature properties of

the BH system. We note here that the quantum phase transition exists in the strict sense only at $T = 0$. For finite T the compressibility, i.e. the change of the particle number per site with the chemical potential, is always nonzero and thus only an approximate Mott phase exists. Furthermore the transition between superfluid and insulating behaviour passes through an intermediate thermal region. Recently this thermal crossover has been studied within a slave-boson formalism [23]. In this work we study the finite- T properties using a perturbative analytic approach in the strong-coupling limit, deriving simple analytical expressions for the occupation number and number fluctuations which cover both quantum and thermal effects. To verify these results numerically in one spatial dimension, we employ a finite- T version of the DMRG approach [21]. We observe good agreement between perturbative and DMRG results, including the description of the thermal crossover between the Mott and insulating phases.

We consider a d -dimensional cubic lattice of N nonlinear oscillators characterised by the following Hamiltonian,

$$\hat{\mathcal{H}} - \mu \hat{\mathcal{N}} = \hat{\mathcal{H}}_{\text{hop}} + \hat{\mathcal{H}}_0, \quad (1)$$

where μ is the chemical potential, $\hat{\mathcal{N}} = \sum_{\mathbf{k}} \hat{n}_{\mathbf{k}}$ is the operator of the total number of atoms, and $\hat{n}_{\mathbf{k}} = \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}$, with $\hat{a}_{\mathbf{k}}^\dagger, \hat{a}_{\mathbf{k}}$ being the usual bosonic creation/annihilation pairs. The lattice sites are numbered with a d -dimensional index $\mathbf{k} = \{k_1, \dots, k_d\}$, while $\sum_{\mathbf{k}}$ means summation over all lattice sites. The first term,

$$\hat{\mathcal{H}}_{\text{hop}} = -J \sum_{\langle \mathbf{k} \mathbf{l} \rangle} \left(\hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{l}} + \hat{a}_{\mathbf{l}}^\dagger \hat{a}_{\mathbf{k}} \right), \quad (2)$$

on the RHS of (1) is the hopping Hamiltonian, $J \geq 0$; the sum extends over nearest neighbours and we assume

cyclic boundary conditions. The second term on the RHS of (1), $\hat{H}_0 = \sum_{\mathbf{k}} H_{\text{nl}}(\hat{n}_{\mathbf{k}})$, is a sum of the nonlinear site Hamiltonians

$$\begin{aligned} H_{\text{nl}}(\hat{n}_{\mathbf{k}}) &= \frac{U}{2} \hat{n}_{\mathbf{k}} (\hat{n}_{\mathbf{k}} - 1) - \mu \hat{n}_{\mathbf{k}} \\ &= \frac{U}{2} (\hat{n}_{\mathbf{k}} - \bar{n})^2 + \text{const}, \end{aligned} \quad (3)$$

where $\bar{n} = \mu/U + 1/2$ is the filling. The quantum averaging is defined in the standard manner as $\langle \dots \rangle = Z^{-1} \text{Tr} [e^{-\beta(\hat{H} - \mu\hat{N})}(\dots)]$, where $Z = \text{Tr} e^{-\beta(\hat{H} - \mu\hat{N})}$ is the statistical sum, and $\beta = 1/T$ is the inverse temperature. Note that oscillator units with $\hbar = k_B = 1$ are used throughout this paper.

While elaborate methods are needed if we wish to exactly pinpoint lobe boundaries or calculate long-range correlations, local quantities like the average number of atoms and the number fluctuations may be obtained at lesser cost. Let us firstly consider the Mott insulator phase. To zero order in hopping, the ground state of the Mott insulator is $|0\rangle_{J=0} = \prod_{\mathbf{k}} |n_0\rangle_{\mathbf{k}}$, where $n_0 = \text{round}(\bar{n})$, cf. Eq. (3). A first order perturbative correction to this state will contain all states which are found from $|0\rangle_{J=0}$ by moving one atom to a neighboring site. All such “single-hop” states are eigenstates of $\mathcal{H}_0$ with the same relative energy U , resulting in an especially simple expression for the ground state to first order in J [24]. Up to second order we find

$$|0\rangle = \alpha \left(\mathbb{1} - \frac{1}{U} \mathcal{H}_{\text{hop}} \right) |0\rangle_{J=0} + \mathcal{O}(\mathcal{H}_{\text{hop}}^2), \quad (4)$$

where α is a normalisation factor. By direct calculation we then find

$$\begin{aligned} \langle 0 | \hat{n}_{\mathbf{k}} | 0 \rangle &= n_0, \\ \delta n^2 &= \langle 0 | \hat{n}_{\mathbf{k}}^2 | 0 \rangle - \langle 0 | \hat{n}_{\mathbf{k}} | 0 \rangle^2 = \frac{2zJ^2 n_0(n_0 + 1)}{U^2}, \end{aligned} \quad (5)$$

where $z = 2d$ is the number of nearest neighbours in the lattice. These formulae are expected to be a good approximation deep in the Mott insulator phase where a large energy gap exists and the thermal occupation of higher levels can be disregarded.

It is worth noting that, if staying within the perturbation approach, these expressions cannot be extended to the thermodynamic limit $N \rightarrow \infty$. To demonstrate this, consider the formula thus obtained for the normalisation constant, α :

$$\begin{aligned} \alpha^2 \left(1 + \frac{1}{U^2} \langle 0 | \mathcal{H}_{\text{hop}}^2 | 0 \rangle \right) &= 1, \\ \langle 0 | \mathcal{H}_{\text{hop}}^2 | 0 \rangle &= zJ^2 N n_0(n_0 + 1). \end{aligned} \quad (6)$$

These results hold only if $zJ^2 N n_0(n_0 + 1) \ll U^2$, therefore making the thermodynamic limit impossible. (This problem went unnoticed in Ref. [24].) On the other hand, local (on-site) quantities should become independent of

the size of the system when it becomes large enough (thermodynamic limit). In fact, our DMRG simulations show that Eq. (5) equally holds if $zJ^2 N n_0(n_0 + 1) \gtrsim U^2$.

In the superfluid phase, one has to account for states with the total number of quanta different from $N n_0$. In the limit of zero hopping an arbitrary eigenstate is a product state, $|\vec{n}\rangle = \prod_{\mathbf{k}} |n_{\mathbf{k}}\rangle_{\mathbf{k}}$. With $\bar{n}$ close to a half-integer, it suffices to keep only two states for each site, $|n_0\rangle$ and $|n_0 + 1\rangle$, effectively turning bosons into fermions. We shall call the subspace of states with $n_{\mathbf{k}} = n_0, n_0 + 1$, the f -subspace. In this subspace we introduce standard fermionic creation and annihilation operators, $\hat{c}_{\mathbf{k}} \hat{c}_{\mathbf{l}}^\dagger + \hat{c}_{\mathbf{l}}^\dagger \hat{c}_{\mathbf{k}} = \delta_{\mathbf{k}\mathbf{l}}$. Within the f -subspace, $(l, m \geq 0)$

$$\begin{aligned} (\hat{a}_{\mathbf{k}}^\dagger)^{l+m} \hat{a}_{\mathbf{k}}^l &= (c_{\mathbf{k}}^\dagger \sqrt{n_0 + 1})^m (n_0 + \hat{c}_{\mathbf{k}}^\dagger \hat{c}_{\mathbf{k}})^l, \\ (\hat{a}_{\mathbf{k}}^\dagger)^l \hat{a}_{\mathbf{k}}^{l+m} &= (n_0 + \hat{c}_{\mathbf{k}}^\dagger \hat{c}_{\mathbf{k}})^l (c_{\mathbf{k}} \sqrt{n_0 + 1})^m. \end{aligned} \quad (7)$$

Using these relations one can express any bosonic operator in fermionic terms. In particular,

$$H_{\text{nl}} = \frac{U}{2} (\hat{c}_{\mathbf{k}}^\dagger \hat{c}_{\mathbf{k}} + n_0 - \bar{n})^2 = \Delta E \hat{c}_{\mathbf{k}}^\dagger \hat{c}_{\mathbf{k}} + \text{const}, \quad (8)$$

where $\Delta E = U(1/2 + n_0 - \bar{n})$. Projection onto the f -subspace thus makes the Hamiltonian linear so it can be directly diagonalised: (omitting an additive constant)

$$\hat{\mathcal{H}} - \mu\hat{N} = \sum_{\mathbf{l}} (E_{\varphi_{l_1}} + \dots + E_{\varphi_{l_d}}) \hat{f}_{\mathbf{l}}^\dagger \hat{f}_{\mathbf{l}}, \quad (9)$$

where $\varphi_l = 2\pi(l-1)/N$, $E_\varphi = \Delta E - 2J(n_0 + 1) \cos \varphi$, and $\hat{f}_{\mathbf{l}}$ are related to $\hat{c}_{\mathbf{k}}$ by a d -dimensional discrete Fourier transform. The average on-site particle number and number fluctuations are then expressed by the average number of “fermions” per site, $\Delta n = \langle \hat{c}_{\mathbf{k}}^\dagger \hat{c}_{\mathbf{k}} \rangle$, as

$$\begin{aligned} \langle \hat{n}_{\mathbf{k}} \rangle &= n_0 + \Delta n, \\ \langle \hat{n}_{\mathbf{k}}^2 \rangle &= \left\langle \left(n_0 + \hat{c}_{\mathbf{k}}^\dagger \hat{c}_{\mathbf{k}} \right)^2 \right\rangle = n_0^2 + (2n_0 + 1)\Delta n, \\ \delta n^2 &= \langle \hat{n}_{\mathbf{k}}^2 \rangle - \langle \hat{n}_{\mathbf{k}} \rangle^2 = \Delta n(1 - \Delta n). \end{aligned} \quad (10)$$

For Δn we readily find: ($\beta = 1/T$)

$$\Delta n = \left(\prod_{s=1}^d \int_0^{2\pi} \frac{d\varphi_s}{2\pi} \right) \frac{1}{1 + \exp \beta \sum_{s=1}^d E_{\varphi_s}}. \quad (11)$$

This formula holds under the thermodynamic limit $N \rightarrow \infty$.

The relations (5) and (10), (11) are found under mutually exclusive conditions. Equation (5) follows if we assume that only the ground state matters, and retain the first non-vanishing correction to its wave function; with only one level being important temperature is of no concern. Conversely, equations (10), (11) ignore corrections to the eigenstate wavefunctions yet account for energy shifts; with many levels being accounted for, thermal

properties are retained. The question we now address is whether a relation can be derived which covers both insulator and superfluid regions, plus, even more importantly, the crossover region (which is naturally termed the thermal region)? It would be natural to include perturbative corrections to all “fermionic” states, but this results in some intractable algebra. It turns out, however, that for $J \ll U$, simple interpolating formulae between the two above results may be found which apply not only to the insulator and superfluid regions, but also to the thermal crossover region between them. Namely,

$$\begin{aligned} \langle \hat{n}_{\mathbf{k}} \rangle &= n_0 + \Delta n, \\ \delta n^2 &= \Delta n(1 - \Delta n) + \frac{2zJ^2(n_0 + \Delta n)(n_0 + \Delta n + 1)}{U^2}, \end{aligned} \quad (12)$$

where Δn is given by Eq. (11). The idea behind (12) is relatively simple: for small J , the quantum contribution (5) is negligible in the thermal and (even more so) the superfluid regions, while in the insulator region, $\Delta n \approx 0$ (or 1), so that (12) coincide with (5).

To verify these results in one spatial dimension, we use a finite-temperature version of the DMRG approach [21]. At the core of our approach is the following block-doubling algorithm: assuming that we know the optimised states for a block of length L , we can then find eigenstates of a ring of three such blocks, (recall that we use periodic boundary conditions) and calculate the thermal ρ -matrix. Tracing over the states of one block yields the ρ -matrix of a double-sized block. Diagonalising this ρ -matrix and taking a certain number of eigenstates corresponding to the largest eigenvalues results in the optimised basis in a block of length $2L$. The loss of probability is assessed by summing the neglected eigenvalues; if this loss becomes unacceptable the iterations are stopped. To initiate the algorithm, we start from a ring of $L_0 + L'$ oscillators, and then trace out the states of L' oscillators, resulting in an optimised basis for a block of L_0 sites. We found this to yield much better results than starting from an open-ended block by simply selecting its lowest eigenstates. The full details of the algorithm will be discussed elsewhere.

A word of caution is necessary here. While borrowing much of the technical side of the DMRG method, our approach is conceptually different, the difference residing in the importance of temperature. Similar to other renormalisation-group methods, DMRG relies on the fact that the dimension of the physically relevant subspace of the full Hilbert space does not grow with the system size. For finite temperatures this can only be true until the

thermal correlation length is reached. Consequently we only grow the block up to a size comparable to the thermal correlation length, and must simply stop as soon as the probability loss becomes unacceptable. We use periodic rather than open boundary conditions based on the same argument. For the temperatures considered here the maximum block length reached in this way is long enough such that finite-size effects are unimportant. For higher temperatures a combination of stochastic and DMRG techniques can be used, which we are developing and will be discussed in a later work.

The results of our calculations using both DMRG and perturbative approaches in 1-D are summarised in the figure, where we plot average on-site numbers and number fluctuations for three “cross-sections” along the μ -axis, for $J = 0.01U$, $0.02U$, $0.05U$. Each plot shows data for two temperatures, $T = 0.01U$ and $T = 0.001U$. The top and bottom rows of plots represent, respectively, the on-site population, $\langle \hat{n}_{\mathbf{k}} \rangle$, and number fluctuations, $\delta n^2 = \langle \hat{n}_{\mathbf{k}}^2 \rangle - \langle \hat{n}_{\mathbf{k}} \rangle^2$. To gain more insight into the thermal region, the middle row of plots shows the difference, Δn , between the on-site population and the nearest integer value. At $T = 0.001U$, the insulator region is clearly defined by Δn abruptly falling to zero. At $T = 0.01U$, Δn is a smooth function vanishing as μ goes deeper into the insulator region. A similar effect is seen for the on-site number fluctuations. At $T = 0.001U$, the phase transition points are well defined, whereas at $T = 0.01U$ the boundaries of the insulator phase are “eroded” so that δn^2 is a visually smooth function of μ .

Comparing the DMRG to perturbative results, we see that the latter provide a surprisingly good description of the quantities in question. For $J = 0.01U$ (left column of plots), we find a very good agreement between the perturbative and DMRG results. This agreement is reasonable for $J = 0.02U$, (center column) becoming only fair at $J = 0.05U$ (right column). Note that, even in the latter case, the error is mostly in positioning the thermal region between the insulator and superfluid phases. Away from the thermal region, the results of the perturbative approach show good agreement with the DMRG results, both for the superfluid and for the insulator.

Acknowledgments

This research was supported by the Deutsche Forschungsgemeinschaft under contract FL210/12 and the New Zealand Foundation for Research, Science and Technology (Grant No. UFRJ0001).

-
- [1] Matthew P. A. Fisher, *Phys. Rev. B* **89**, 546 (1989).
 - [2] Subir Sachdev, *Quantum Phase Transitions* (Cambridge University Press, 1999).
 - [3] D. Jaksch *et al.*, *Phys. Rev. Lett.* **81**, 3108 (1998).

- [4] M. Greiner *et al.*, *Nature*, **415**, 39 (2002).
- [5] Olaf Mandel *et al.*, e-print cond-mat/0301169
- [6] D.Jaksch *et al.*, *Phys. Rev. Lett.* **82**, 1975 (1999); G.Brennen *et al.*, *Phys. Rev. Lett.* **82**, 1060 (1999).




FIG. 1: Average number and number fluctuations *vs* μ , for two temperatures ($T = 0.01U$, $0.001U$) and three values of the hopping parameter ($J = 0.01U$, $0.02U$, $0.05U$). Top row: on-site population, $n = \langle \hat{n}_k \rangle$; middle row: difference, Δn , between $\langle \hat{n} \rangle$ and the nearest integer; bottom row: on-site number fluctuations, $\delta n^2 = \langle \hat{n}_k^2 \rangle - \langle \hat{n}_k \rangle^2$. The lines show perturbative results ($T = 0.001U$: solid line, $T = 0.01U$: dashed line); the markers show results of DMRG calculations ($T = 0.001U$: diamonds, $T = 0.01U$: circles). Open markers are used for $\Delta n < 0$.

- [7] P. Bouyer, M.A. Kasevich, Phys. Rev. **A** **56**, R1083 (1997).
- [8] P. Rabl *et al.*, e-print cond-mat/0304026.
- [9] Daniel S. Rokhsar and B. G. Kotliar, Phys. Rev. B **44**, 10328 (1991).
- [10] J. K. Freericks, and H. Monien, Europhys. Lett., **26**, 545 (1994).
- [11] J. K. Freericks, and H. Monien, Phys. Rev. B **53**, 2691 (1996).
- [12] N. Elstner, and H. Monien, Phys. Rev. B **59**, 12184 (1999).
- [13] N. Elstner, and H. Monien, *cond-mat/9905367* (1999).
- [14] W. Krauth and N. Trivedi, Europhys. Lett. **14**, 627 (1991).
- [15] G.G. Batrouni, R.T. Scalettar, and G.T. Zimanyi, Phys. Rev. Lett. **65**, 1765 (1990).
- [16] R. T. Scalettar *et al.*, Phys. Rev. B **51**, 8467 (1995).
- [17] G. G. Batrouni *et al.*, Phys. Rev. Lett. **89**, 117203 (2002).
- [18] V. A. Kashurnikov, N. V. Prokof'ev, and B. V. Svistunov, Phys. Rev. A **66**, 031601 (2002).
- [19] L. Amico and V. Penna, Phys. Rev. Lett. **80**, 2189-2192 (1998); Phys. Rev. **B** **62**, 1224-1237 (2000).
- [20] D. van Oosten, P. van der Straten, H. T. C. Stoof, Phys. Rev. A **63**, 053601 (2001).
- [21] S. R. White, Phys. Rev. B **48**, 10345 (1993).
- [22] Till D. Kühner, Steven R. White, and H. Monien, Phys. Rev. B **61**, 12474 (2000).
- [23] D.B.M. Dickerscheid *et al.*, e-print cond-mat/0306573.
- [24] D. C. Roberts and K. Burnett, Phys. Rev. Lett. **90**, 150401 (2003).